

Death Receptor 3 promotes chemokine directed leukocyte recruitment in acute resolving inflammation and is essential for pathological development of mesothelial fibrosis in chronic disease

William Victor Perks^{*‡}, Ravinder Kaur Singh^{*‡}, Gareth Wyn Jones[†], Jason Peter Twohig^{*†}, Anwen Sian Williams[†], Ian Robert Humphreys^{*}, Philip Russel Taylor[†], Simon Arnett Jones[†], Eddie Chung Yern Wang^{*§}

Departments of ^{*}Medical Microbiology & Infectious Disease and [†]Rheumatology, Cardiff Institute of Infection & Immunity, School of Medicine, Cardiff University, Heath Park, Cardiff CF14 4XN, UK

[‡] These authors contributed equally to the manuscript

No of pages: 25 (including References and Figure legends); **4 Figures**

Sources of support: WVP was supported by a PhD studentship from the I3-IRG, joint funded by the British Medical Research Council (MRC) and the School of Medicine, Cardiff University awarded to ECYW and SAJ. JPT was employed on MRC grant G0901119 awarded to ECYW and IRH. RKS was funded by BBSRC/GSK CASE studentship BB/H530589/1.

[§]**Corresponding author:** Dr. E.C.Y. Wang, Department of Medical Microbiology & Infectious Disease, Cardiff Institute of Infection & Immunity, School of Medicine, Cardiff University, Henry Wellcome Building, Heath Park, Cardiff CF14 4XN, UK

e-mail: wangec@cf.ac.uk

ABSTRACT

Death Receptor 3 (DR3, TNFRSF25) and its TNF-like ligand TL1A (TNFSF15) control a number of processes in inflammatory diseases through the expansion of effector T cells and the induction of pro-inflammatory cytokines from myeloid and innate lymphoid cells. Here, using wildtype (DR3^{+/+}) and DR3 knockout (DR3^{-/-}) mice, we show that the DR3/TL1A pathway triggers the release of multiple chemokines following acute peritoneal inflammation initiated by a single application of *Staphylococcus epidermidis* supernatant (SES), correlating with the infiltration of multiple leukocyte subsets. In contrast, leukocyte infiltration was not DR3-dependent following viral challenge with murine cytomegalovirus (MCMV). DR3 expression was recorded on connective tissue stroma, which provided DR3-dependent release of CCL2 (MCP-1), CCL7 (MCP-3), CXCL1 (KC), and CXCL13 (BCA-1). CCL3 (MIP-1 α), CCL4 (MIP-1 β) and CXCL10 (IP10) production was also DR3-dependent, but qRT-PCR showed that their derivation was not stromal. *In vitro* cultures identified resident macrophages as a DR3-dependent source of CCL3. Whether DR3 signaling could contribute to a related peritoneal pathology was then tested using multiple applications of SES, the repetitive inflammatory episodes of which lead to peritoneal membrane thickening and collagen deposition. Unlike their DR3^{+/+} counterparts, DR3^{-/-} mice did not develop fibrosis of the mesothelial layer. Thus, this work describes both a novel function and essential requirement for the DR3/TL1A pathway in acute, resolving and chronic inflammation in the peritoneal cavity.

1 INTRODUCTION

2 Death Receptor 3 (DR3, TRAMP, LARD, WSL-1, Apo-3, TR3, TNFRSF25)¹⁻⁶, a member of the
3 tumor necrosis factor receptor superfamily (TNFRSF), and its TNFSF ligand TL1A (TNFSF15)⁷,
4 have emerged as major regulators of inflammation and immunity. Genome-wide association studies
5 (GWAS) have consistently linked polymorphisms around the *TNFSF15* gene as a risk factor in
6 inflammatory bowel diseases (IBD) such as ulcerative colitis, Crohn's disease and/or irritable bowel
7 syndrome⁸⁻¹¹, and in the arthritic disease ankylosing spondylitis¹², with associations also reported
8 with primary billiary cirrhosis¹³. Furthermore, variation in the *TNFRSF25* gene locus has been
9 suggested as a risk factor in rheumatoid arthritis (RA)¹⁴, while other genetic studies have associated
10 the *TNFSF15* gene with leprosy¹⁵, implicating a further role for the DR3/TL1A axis in pathogenic
11 immunity.

12
13 These wide-ranging links to multiple pathologies have been supported by *in vitro* and *in vivo* studies
14 in both man and mouse. Levels of serum and tissue TL1A are increased in RA¹⁶, IBD¹⁷, psoriasis¹⁸,
15 primary billiary cirrhosis¹⁹ and ankylosing spondylitis²⁰, while mice overexpressing TL1A develop
16 colitis or ileitis²¹⁻²³. In contrast, mice deficient for DR3 or TL1A, or those expressing dominant
17 negative forms of DR3 are resistant to multiple models of inflammatory and autoimmune disease²⁴⁻
18 ³¹, or exhibit impaired immunity to bacterial or viral pathogens^{32, 33} with a more debated impact on
19 immunity to parasitic worms^{34, 35}. Much of DR3 function has been attributed to its expression on T
20 cells (CD4⁺ regardless of lineage or CD8⁺) and the capacity of TL1A, either directly or indirectly, to
21 drive the proliferation of pathogenic effector T cell numbers^{26, 27, 36, 37}. A single study also suggests a
22 role in inhibiting proliferation of B cells³⁸. This has defined significant roles for the DR3/TL1A
23 pathway in adaptive immune responses.

24
25 However, DR3 expression is also found on innate immune cell and non-hematopoietic lineages. DR3
26 signaling on myeloid cells influences myeloid differentiation^{30, 39}, while on NKT and innate

lymphoid cells (ILCs), triggers effector cytokine release and expansion essential for the development of airway inflammation^{25, 34}. Further, DR3^{-/-} mice exhibit behavioural defects associated with impaired corticostriatal innervation controlled by DR3 expression on neurones⁴⁰ and its expression on tubular epithelial cells regulates renal inflammation and injury caused by reperfusion^{41, 42}. On osteoblasts, DR3 can trigger apoptosis *in vitro*⁴³ or control bone mineral apposition *in vivo*⁴⁴. These findings indicate that the DR3/TL1A pathway has intrinsic functions and also controls early events in immune processes in addition to the more established roles in acquired immunity.

In this report, we further investigated the *in vivo* role of the DR3/TL1A pathway using sterile challenge of *Staphylococcus epidermidis* supernatant (SES) in the peritoneal cavity^{45, 46}. As a single application, this system mimics a resolving inflammatory response that returns to baseline levels⁴⁵, allowing scrutiny of infiltrating cells and soluble factors in peritoneal washouts within hours of the challenge. Using multiple applications, pathology develops in the form of peritoneal membrane fibrosis⁴⁷, mirroring the pathology associated with repeated bacterial infections in renal patients on peritoneal dialysis. Consequently, we show that the DR3/TL1A axis shapes the inflammatory response within the peritoneum and report a previously unappreciated role for DR3 in controlling inflammation via modulation of chemokine expression by both hematopoietic and stromal cells.

MATERIALS AND METHODS

Animals

Age-matched DR3^{-/-} and DR3^{+/+} littermates of 8-12 weeks of age on a C57BL/6 background were used in experiments, derived from a DR3^{het} colony that was founded from mice provided by CRUK⁴⁸. For peritoneal challenge with SES, animals of mixed sex were used as no significant differences in peritoneal leukocyte content of male and female mice were observed, either at baseline or during the course of experiments (data not shown). Irrespective, the male:female ratio in each DR3^{+/+} and DR3^{-/-} group for each timepoint were matched. For challenge with MCMV, all animals used were male. All procedures were approved by the Local Research Ethics Committee and performed in accordance with Home Office approved license PPLs 30/2401 (for SES), 30/2442 and 30/2580 (for MCMV).

Immunohistochemistry

Briefly, peritoneal membranes were harvested and fixed in neutral buffered formal saline and permeated with paraffin wax before 5µm sectioning. Sections were rehydrated and endogenous peroxidase activity was blocked. Antigen unmasking was achieved by incubating the sections in Proteinase K (20 µg/ml in TE buffer; Sigma) for 10 mins. Following blocking steps (20% goat serum and streptavidin/biotin blocking kit (Vector Laboratories), sections were incubated overnight with 4 µg/ml goat biotinylated anti-DR3 (R&D systems) or isotype controls diluted in PBS followed by streptavidin-HRPO secondary antibody (Vector Laboratories) as per manufacturer's instructions. Positive staining was visualized using a DAB chromogen (Vector Laboratories). Collagen was visualized using van Gieson stain. Sections were counterstained with hematoxylin, dehydrated and mounted in DPX. Images were captured using an Olympus N457 or Canon 100D digital camera and positive staining analyzed using Adobe Photoshop CS3.5. Randomly selected areas were used for analysis with the average from 5 being used to generate values for each section.

qRT-PCR

Short title: DR3 regulates peritoneal inflammation

Total RNA was extracted following manufacturer's instructions using a RNeasy mini kit (Qiagen, Crawley, UK) from snap frozen and powdered peritoneal membranes or pelleted macrophages. cDNA synthesis and PCR was performed using systems and primers generated through Primer Design (Southampton, UK) on an iCycler qPCR machine (Biorad, Hemel Hempstead, UK) with β -actin as the housekeeping control gene. Relative Quantity (RQ) was calculated after subtracting the negative control. Primers used (forward first, reverse second) were as follows:

CCL2 (5'-CCCAATGAGTAGGCTGGAGA-3' and 5'-TCTGGACCCATTCCTTCTTG-3');
CCL3 (5'-TTTTGAAACCAGCAGCCTTT-3' and 5'-CTGCCTCCAAGACTCTCAGG-3');
CCL4 (5'-CCCCTTCCTGCTGTTTCTC-3' and 5'-CTCACTGGGGTTAGCACAGA-3');
CCL5 (5'-GTGCCCACGTCAAGGAGTAT-3' and 5'-CCACTTCTTCTCTGGGTTGG-3');
CCL7 (5'-CCCAAGAGGAATCTCAAGAGC-3' and 5'-ATAGCCTCCTCGACCCACTT-3');
CXCL1 (5'-GCACCCAAACCGAAGTCATA-3' and 5'-AGGTGCCATCAGAGCAGTCT-3');
CXCL2 (5'-AAAGTTTGCCTTGACCCTGA-3' and 5'-CTTTGGTTCTTCCGTTGAGG-3');
CXCL5 (5'-GCCCTACGGTGGAAAGTCATA-3' and 5'-GTGCATTCCGCTTAGCTTTC-3');
CXCL10 (5'-TGAATCCGGAATCTAAGACCA-3' and 5'-GAGGCTCTCTGCTGTCCATC-3');
CXCL13 (5'-TCGTGCCAAATGGTTACAAA-3' and 5'-GGTGCAGGTGTGTCTTTTGA-3');
TL1A (5'-CAGCAGAAGGATGGCAGA-3' and 5'-CTCTGGCCTGTGTCTACA-3').

Preparation of SES, determination of biological activity and the SES model of peritoneal challenge

SES was prepared and used in peritoneal challenge as described previously^{46, 47}. Supernatants were filtered to remove live bacteria, lyophilized and stored at -70°C. Biological activity was determined by its capacity to induce IL6 release from RAW 264 cells in a 24 hr culture, measured by ELISA (BD Pharmingen OpEIA; BD Biosciences, Oxford, UK).

Challenge with MCMV

Short title: DR3 regulates peritoneal inflammation

Mice were challenged intraperitoneally (i.p.) with 5×10^4 plaque-forming units of MCMV Smith strain (American Type Culture Collection, Manassa, VA, USA), as previously described⁴⁹.

Flow cytometry

Following peritoneal lavage and centrifugation to isolate leukocytes, samples were treated with Fc block (BD Pharmingen) and then stained at 4°C for 25 mins with combinations of the following pre-conjugated mAbs: CD19-FITC, CD8a-FITC, NK1.1-PE, Gr-1-PerCP-Cy5.5, CD4-PerCP-Cy5.5, CD11c-PE-Cy7, Ly6G-V450, CD44-V450, CD11b-APC-Cy7, CD3-APC-Cy7 (all BD Biosciences); B220-PE, F4/80-APC (Invitrogen); Ly6B.2-PE (AbD Serotec); TCR $\alpha\beta$ -APC (Caltag). Samples were then washed, fixed in 1% paraformaldehyde before running and analysis on a CyAn™ ADP Flow Cytometer using Summit software (Beckman Coulter). Cell subsets were identified using the following criteria: neutrophils (7/4⁺, Ly6G⁺), resident macrophages (F4/80⁺CD11b⁺), inflammatory macrophages (F4/80^{int}CD11b^{int}), eosinophils (F4/80^{int}CD11b^{int}SSC^{hi}), NK cells (NK1.1⁺ $\alpha\beta$ TCR⁻), NKT cells (NK1.1⁺ $\alpha\beta$ TCR⁺), T cells (CD3⁺ $\alpha\beta$ TCR⁺) and B cells (CD19⁺B220⁺).

ELISAs and chemokine arrays

ELISAs were performed following manufacturers' instructions. For murine CCL2, CCL3, CCL5, CCL5, CXCL1, CXCL2, CXCL5, CXCL10 and CXCL13, ELISAs were obtained from R&D Systems (Abingdon, UK). ELISA for murine CCL7 was obtained from Antigenix America (NY 11746, USA). For determining concentration of chemokines in cultures of resident macrophages, the Legendplex™ mouse proinflammatory chemokine panel was used following manufacturer's instructions (Biolegend).

Statistical analysis

Staining readouts were percentages, therefore, non-parametric Mann-Whitney U tests were used for statistical analysis. For protein concentrations from ELISAs and RQ from qRT-PCR, student's t-

Short title: DR3 regulates peritoneal inflammation

- 1 Tests and ANOVAs were used. Analyses were performed on GraphPad Prism - p values of <0.05
- 2 were considered significant and values of <0.01 were considered highly significant.
- 3

RESULTS

DR3 and TL1A expression in the peritoneal cavity following SES challenge

The DR3/TL1A pathway has been shown to be essential in the development of a number of models of disease involving intraperitoneal priming or challenge^{25-27, 30, 32, 33}, but little has been reported about DR3 and TL1A expression patterns in the peritoneal tissue and cavity during this process. DR3 expression in the mesothelial layer of the peritoneal membrane was first investigated, before and after SES challenge by staining of sections using a polyclonal antibody. Anti-DR3 showed minimal staining on DR3^{-/-} peritoneal mesothelial layers ($0.25 \pm 0.08\%$ positive pixels), but a significant signal was recorded from DR3^{+/+} samples, in the absence of ($6.5 \pm 0.3\%$) and up to 12 hrs post-challenge ($7.6 \pm 1.6\%$) before becoming negative after 24 hrs ($0.08 \pm 0.04\%$) (Fig. 1A, B). DR3 was also found on CD4⁺, CD8⁺ and NKT cells within the peritoneal cavity, in agreement with previous reports studying DR3 expression in splenocytes (^{25, 33} and data not shown). This data shows for the first time that DR3 is present on the stroma of the peritoneal cavity. In addition, expression of TL1A mRNA was measured using qRT-PCR on peritoneal membrane sections, as well as from resident peritoneal macrophages isolated by cell sorting (F4/80⁺CD11b⁺) and treated *in vitro* with SES. Whilst the mesothelial layers were negative for TL1A throughout the timecourse (Fig. 1C), TL1A mRNA was triggered from resident peritoneal macrophages within 1 hr of SES treatment, irrespective of their derivation from DR3^{+/+} or DR3^{-/-} mice (Fig. 1D). Thus, bacterial challenge in the peritoneal cavity primarily induces TL1A from resident macrophages, not stroma, but responses to TL1A can come from DR3 on both stroma and leukocytes.

Leukocyte accumulation is impaired in the absence of DR3

Leukocyte infiltration of the peritoneal cavity following SES challenge was next examined in the presence or absence of DR3. The composition of inflammatory cells in the peritoneal lavage was assessed using multiparameter flow cytometry and 2 panels of mAbs defining myeloid lineages and lymphoid populations (see Materials and Methods). Before challenge, there were no differences in

the leukocyte content of DR3^{-/-} and DR3^{+/+} peripheral blood or peritoneal lavage (Supplemental Table 1), showing that the absence of DR3 did not influence the baseline cellular constitution of the circulation or peritoneal cavity. The progression of this model from inflammatory insult to its resolution has been well-characterized, demonstrating a biphasic cellular infiltration predominantly consisting of neutrophils that peaks in the first 6 hrs and declines thereafter. Increasing numbers of other cell types follows, peaking after 48 hrs, before falling away over 96 hrs (^{45, 46} and Fig. 2). In DR3^{-/-} mice, significantly lower numbers of neutrophils in the early stage were observed following SES challenge compared to their DR3^{+/+} counterparts. Interestingly, viral challenge with murine cytomegalovirus (MCMV) induced considerably less neutrophil accumulation and showed no differences between DR3^{+/+} and DR3^{-/-} mice (Fig 2A) , consistent with reports that neutrophil recruitment following MCMV challenge is IL-22 dependent ⁵⁰. During the second phase after SES challenge, eosinophils, inflammatory macrophages, T cells, B cells and NKT cells were decreased in number (Fig 2B). While the profile of resident macrophages and NK cell numbers were not significantly different between DR3^{+/+} and DR3^{-/-} mice throughout the time course, peritoneal B cell numbers remained significantly higher in DR3^{+/+} animals by 96 hrs post-challenge (Fig. 2B). Overall, there was an intrinsic defect in the accumulation of leukocytes in DR3^{-/-} mice after SES challenge in the peritoneal cavity.

Proliferation is unaltered in leukocytes from DR3^{-/-} mice

Reduced numbers of effector cells, normally at the site of pathology, have been reported in the absence of DR3 in a variety of different pathological settings, primarily being ascribed to defects in proliferation ^{25-27, 32-34}. We, therefore, measured the levels of proliferation of cells from peritoneal lavage using detection of Ki67 at 48 hrs post-SES challenge, when there were peak numbers of infiltrating leukocytes. Surprisingly, there were no differences in Ki67 staining of any of the cell subsets when comparing DR3^{+/+} and DR3^{-/-} mice (Supplemental Fig. 1 and Supplemental Table 2), indicating that at this early stage of the inflammatory process, DR3 was regulating a distinct

Short title: DR3 regulates peritoneal inflammation

mechanism of cellular accumulation. Because DR3 contains a death-inducing intracellular domain, apoptosis using Annexin-V staining was also measured, but consistent with other reports^{33, 51}, the absence of DR3 did not alter the levels of cell death in peritoneal leukocytes (data not shown). The DR3/TL1A pathway therefore regulated peritoneal leukocyte numbers early after bacterial challenge with a mechanism that was independent of proliferation and cell death.

Levels of multiple chemokines are impaired in DR3^{-/-} mice following SES challenge

Another possible explanation for the defect in accumulation of cellular infiltrate is that DR3 regulates leukocyte recruitment. To ensure that the absence of DR3 did not adversely influence baseline levels, cell-free supernatant from peritoneal lavages of unchallenged DR3^{+/+} and DR3^{-/-} mice were tested for a range of chemokines capable of attracting virtually every subset of leukocytes between them. No significant differences were found at baseline, with detectable levels of CCL3, CCL4, CXCL5 (targeting neutrophils, monocytes), CXCL10 (T cells, NK cells) and CXCL13 (B cells), while CCL2, CCL5, CCL7 (dendritic cells, monocytes, T cells), CXCL1 and CXCL2 (neutrophils) were below levels of detection of their specific kits (Supplemental Fig. 2). The kinetics of chemokine release were then investigated in the peritoneal lavage of DR3^{+/+} and DR3^{-/-} mice challenged with SES. While some chemokines including CCL5, CXCL2 and CXCL5 showed no differences between genotypes, the expression of CCL2, CCL3, CCL4, CCL7, CXCL1, CXCL10, CXCL13 were all significantly impaired in DR3^{-/-}, as compared to DR3^{+/+} mice, although not necessarily at the same timepoint as when peak concentrations were observed (Fig. 3A). In DR3^{-/-} mice, CCL2, CCL7 and CXCL1 were significantly reduced at 1 hr; CXCL10 at 3 hrs; CCL3 and CCL4 at 6 hrs; and CXCL13 at 24 hrs.

Select chemokines are differentially produced from the stroma or resident macrophages in a DR3-dependent fashion

Chemokines can be derived from both stromal and hematopoietic cells. To delineate between these potential sources, qRT-PCR for the messages of all of the above chemokines were performed on RNA extracted from DR3^{+/+} and DR3^{-/-} mesothelial layers at timepoints between 1 and 24 hrs following SES challenge. Those showing no significant differences in protein levels between DR3^{-/-} and DR3^{+/+} mice (CCL5, CXCL2 and CXCL5) also showed no significant differences in message. In contrast, of the remaining 7, only CCL2, CCL7, CXCL1 and CXCL13 (attractants of dendritic cells, neutrophils, eosinophils, memory T and B cells) demonstrated a significant reduction in message from DR3^{-/-} compared to DR3^{+/+} mesothelial layers (Fig. 3B), implicating the stroma as a significant DR3-dependent source of these chemokines and suggesting non-stromal cells are the DR3-dependent source of the others. To further investigate this, resident macrophages (the largest innate immune leukocyte population in the unchallenged peritoneal cavity – see Supplemental Table 1) were extracted from DR3^{+/+} and DR3^{-/-} mice, pooled and challenged with SES *in vitro*. Of 13 chemokines studied, only CCL3 and CCL4 were recorded at levels above the detection limit of the array. DR3^{+/+} resident macrophages produced significantly more CCL3, but not CCL4, compared to DR3^{-/-} resident macrophages in response to SES (Fig. 3C). The overall conclusion is that DR3 signaling controls recruitment of multiple leukocyte subsets following bacterial challenge in the peritoneal cavity by a broad release of chemokines, some of which come from stroma, while others are derived from non-stromal cell types including resident macrophages within the peritoneal cavity.

DR3 deficiency protects against pathology of SES-induced chronic inflammation

To test whether DR3 signaling could contribute to pathology in a related model of disease, DR3^{-/-} animals were challenged with multiple serial applications of SES, a system designed to mimic the fibrosis of the peritoneal membrane observed in peritoneal dialysis patients following persistent and regular bacterial infections in the peritoneal cavity⁴⁷. Thickening of the mesothelial layer and increased deposition of collagen was observed in DR3^{+/+} mice, but DR3^{-/-} mice showed resistance to

Short title: DR3 regulates peritoneal inflammation

1 this (Fig. 4), demonstrating an essential requirement for DR3 in the development of this fibrotic
2 pathology.

3

DISCUSSION

In recent years, the DR3/TL1A pathway has emerged as a master regulator of inflammatory and autoimmune disease, as well as a co-ordinator of immune responses to a variety of pathogens. This is in the context of DR3^{-/-} mice showing no impairments in the development of their immune systems, constitution of their immune (or other) organs⁴⁸ or intrinsic extrathymic T cell trafficking²⁶. Much of the *in vivo* research in the field has used models of disease involving antigen priming or pathogen challenge in the peritoneal cavity, and have demonstrated that antigen-specific splenic responses following such challenges are relatively normal in the absence of DR3 indicating that immune priming is intact. Here, we show that the DR3/TL1A pathway controls the very earliest of immune processes during peritoneal challenge, regulating the production of chemokines that draws pro-inflammatory leukocytes into sites of inflammation. This effect is dependent on the type of challenge, with SES, but not MCMV, inducing DR3-dependent leukocyte infiltration. There is the possibility that the DR3/TL1A pathway interacts with specific TLR signaling pathways as *Staphylococcus epidermidis* is recognized through TLR2⁵², while MCMV triggers TLR3 and 9⁵³ and induces neutrophil recruitment through IL-22⁵⁰.

Prior to this report, DR3 function has primarily been attributed to the expression of DR3 on multiple cell types of the hematopoietic lineage, ranging from NKT, CD4⁺, CD8⁺ T and B cells to ILCs, macrophages and neutrophils. This is the first report of significant DR3 expression on a stromal layer in an *in vivo* setting and in the absence of active immune responses. Previous studies have indicated that DR3 is inducible on stroma by pro-inflammatory events, as has been described for tubular epithelial cells following ischemic injury⁴¹ and possibly in joints undergoing antigen-induced arthritis, although the latter could be attributed to infiltrating cells⁵¹. More intriguingly, stromal DR3 expression seems to be closely regulated in the peritoneal cavity, with the signal disappearing 24 hrs after SES challenge. In contrast, although endothelial cells have been reported to release TL1A⁷, its major source following SES challenge was resident macrophages, occurring within an hour of

stimulation and independent of DR3 expression (Fig. 1). This is in agreement with previous reports showing cells of the myeloid lineage respond to bacterial challenges by rapid TL1A release^{26, 32}.

DR3-dependent release of some chemokines has been previously reported, but not on the scale shown here. Thus, *in vitro* studies on the myeloid THP-1 cell line has demonstrated IL-8 release requiring TL1A and IFN γ priming⁵⁴, while CXCL1 is reduced in the joints of DR3^{-/-} mice undergoing AIA⁵¹. Both these chemokines, in humans and mice respectively, are established neutrophil attractants. This report indicates that at least in the murine SES challenge system, the production of CXCL1, but not 2 other neutrophil chemoattractants (CXCL2/MIP-2 and CXCL5/LIX), is DR3-dependent. It also shows, for the first time, that there is a stromal contribution to this (Fig. 3 and 4). Whether this is the dominant DR3-dependent chemokine controlling neutrophil attraction remains an area of further study as there are multiple other chemokine/chemokine receptor interactions capable of this⁵⁵, many of which have not been tested in this study.

The stromal release of a number of other chemokines demonstrated DR3-dependency, namely CCL2, CCL7 and CXCL13. CCL2 and CXCL13 have primary chemokine receptors, CCR2 and CXCR5 respectively. CCR7 is more pleiotropic, binding CCR1, 2 and 3, together having the capacity to attract monocytes, eosinophils, DCs, T and B cells⁵⁵. This correlated well with the subsets of leukocytes that showed impaired SES-induced accumulation in the peritoneal cavity of DR3^{-/-} mice. Interestingly, however, there were other chemokines in DR3^{-/-} peritoneal lavage that were reduced in concentration, namely CCL3, CCL4 and CXCL10. Their ligands include CCR1, CCR5 and CXCR3 found on monocytes, T, NK and mast cells. We further identified resident macrophages as a DR3-dependent source of CCL3 (Fig 3C). The assumption is that CCL4 and CXCL10 were derived from other as yet unidentified leukocytes as these chemokines were either not produced or did not show DR3-dependent production by resident macrophages, while their mRNA levels were unaltered in the DR3^{-/-} mesothelium. Further studies would be required to identify which other cells are producing

DR3-dependent CCL4 and CXCL10, and determine the relative contribution of DR3 on stroma versus resident versus infiltrating leukocytes to recruitment in this system. The very rapid peak of production for many of the chemokines (less than 3 hrs – see Fig 3) would indicate either a stromal or resident leukocyte source for these, with the possibility of infiltrating leukocytes contributing to levels of chemokines at later time points. In this context, DR3 expression on innate immune cells is constitutive on NKT cells (which have been shown to produce IL-4 and IL-13 in models of allergic lung inflammation)²⁵, or can be induced on macrophages as well as neutrophils⁵¹. The latter can also produce multiple chemokines and is a major infiltrating cell type⁵⁶.

The leukocyte recruitment function of the DR3/TL1A pathway shows distinct differences to the role that has been characterized for IL-6 using intraperitoneal SES challenge. While both impact on T cell recruitment correlating with lower levels of CCL4 and CXCL10, IL-6^{-/-} animals display a reduction in CCL5⁵⁷, which was not observed in DR3^{-/-} mice. Furthermore, unlike DR3^{-/-} animals, IL-6^{-/-} mice show normal early neutrophil⁴⁶ and later B cell⁵⁷ infiltration accompanied by unaltered levels of CXCL1 and CXCL13. This indicates that DR3 signaling is distinct and not directly upstream of IL-6. More recently, IL-6 has been shown to be essential for the development of peritoneal membrane fibrosis following multiple acute inflammatory episodes induced by serial applications of SES. This was attributed to IL-6-mediated Th1 effector commitment⁴⁷. Our data indicate that this model is also dependent on DR3, but likely through its general regulation of effector T cells. DR3^{-/-} mice are resistant to the induced increases in peritoneal membrane thickening and collagen deposition. This is consistent with the phenotype of transgenic mice overexpressing TL1A, which spontaneously develop intestinal and colonic fibrosis accompanied by goblet and Paneth cell hyperplasia^{22, 23, 58}. One significant question is the relative contribution to pathology of DR3's role in leukocyte recruitment versus its function in driving effector cell expansion^{22, 26, 27, 29, 33, 34, 59}. While this can be divided into early and late inflammatory processes in the SES challenge model, the two immunological events are not mutually exclusive and could explain some previously observed

1 phenomena, such as the altered localization of T cells and macrophages in DR3^{-/-} mice undergoing
2 OVA-induced lung inflammation²⁶.

3

4 It is evident from this work that DR3/TL1A signaling regulates primary inflammatory responses,
5 with potential impact on downstream adjuvant and host defense activities. The understanding of these
6 roles to pathological processes is an area of further research that will be important for developing the
7 DR3/TL1A pathway as a therapeutic target in inflammatory disease, agonism or antagonism of which
8 may prevent disease depending on the effector T cells being influenced.

9

1 **Acknowledgements**

2 Thanks to the Central Biotechnology Service, School of Medicine, Cardiff University for access to
3 flow cytometers. WVP, GWJ, JPT and RKS performed experiments. ASW, IRH, PRT, SAJ and
4 ECYW contributed through conceptual and experimental design, animal models and writing of the
5 manuscript.

6

7 **Conflict of Interest Disclosure**

8 The authors declare no commercial or financial conflict of interest.

9

REFERENCES

- [1] Bodmer JL, Burns K, Schneider P, Hofmann K, Steiner V, Thome M, Bornand T, Hahne M, Schroter M, Becker K, Wilson A, French LE, Browning JL, MacDonald HR, Tschopp J: TRAMP, a novel apoptosis-mediating receptor with sequence homology to tumor necrosis factor receptor 1 and Fas(Apo-1/CD95). *Immunity* 1997, 6:79-88.
- [2] Chinnaiyan AM, O'Rourke K, Yu GL, Lyons RH, Garg M, Duan DR, Xing L, Gentz R, Ni J, Dixit VM: Signal transduction by DR3, a death domain-containing receptor related to TNFR-1 and CD95. *Science* 1996, 274:990-2.
- [3] Kitson J, Raven T, Jiang YP, Goeddel DV, Giles KM, Pun KT, Grinham CJ, Brown R, Farrow SN: A death-domain-containing receptor that mediates apoptosis. *Nature* 1996, 384:372-5.
- [4] Marsters SA, Sheridan JP, Donahue CJ, Pitti RM, Gray CL, Goddard AD, Bauer KD, Ashkenazi A: Apo-3, a new member of the tumor necrosis factor receptor family, contains a death domain and activates apoptosis and NF-kappa B. *Curr Biol* 1996, 6:1669-76.
- [5] Tan KB, Harrop J, Reddy M, Young P, Terrett J, Emery J, Moore G, Truneh A: Characterization of a novel TNF-like ligand and recently described TNF ligand and TNF receptor superfamily genes and their constitutive and inducible expression in hematopoietic and non-hematopoietic cells. *Gene* 1997, 204:35-46.
- [6] Screaton GR, Xu XN, Olsen AL, Cowper AE, Tan R, McMichael AJ, Bell JI: LARD: a new lymphoid-specific death domain containing receptor regulated by alternative pre-mRNA splicing. *Proc Natl Acad Sci U S A* 1997, 94:4615-9.
- [7] Migone TS, Zhang J, Luo X, Zhuang L, Chen C, Hu B, Hong JS, Perry JW, Chen SF, Zhou JX, Cho YH, Ullrich S, Kanakaraj P, Carrell J, Boyd E, Olsen HS, Hu G, Pukac L, Liu D, Ni J, Kim S, Gentz R, Feng P, Moore PA, Ruben SM, Wei P: TL1A is a TNF-like ligand for DR3 and TR6/DcR3 and functions as a T cell costimulator. *Immunity* 2002, 16:479-92.
- [8] Yamazaki K, McGovern D, Ragoussis J, Paolucci M, Butler H, Jewell D, Cardon L, Takazoe M, Tanaka T, Ichimori T, Saito S, Sekine A, Iida A, Takahashi A, Tsunoda T, Lathrop M, Nakamura Y: Single nucleotide polymorphisms in TNFSF15 confer susceptibility to Crohn's disease. *Hum Mol Genet* 2005, 14:3499-506.
- [9] Jostins L, Ripke S, Weersma RK, Duerr RH, McGovern DP, Hui KY, Lee JC, Schumm LP, Sharma Y, Anderson CA, Essers J, Mitrovic M, Ning K, Cleynen I, Theatre E, Spain SL, Raychaudhuri S, Goyette P, Wei Z, Abraham C, Achkar JP, Ahmad T, Amininejad L, Ananthakrishnan AN, Andersen V, Andrews JM, Baidoo L, Balschun T, Bampton PA, Bitton A, Boucher G, Brand S, Buning C, Cohain A, Cichon S, D'Amato M, De Jong D, Devaney KL, Dubinsky M, Edwards C, Ellinghaus D, Ferguson LR, Franchimont D, Fransen K, Gearry R, Georges M, Gieger C, Glas J, Haritunians T, Hart A, Hawkey C, Hedl M, Hu X, Karlsen TH, Kupcinskis L, Kugathasan S, Latiano A, Laukens D, Lawrance IC, Lees CW, Louis E, Mahy G, Mansfield J, Morgan AR, Mowat C, Newman W, Palmieri O, Ponsioen CY, Potocnik U, Prescott NJ, Regueiro M, Rotter JJ, Russell RK, Sanderson JD, Sans M, Satsangi J, Schreiber S, Simms LA, Sventoraityte J, Targan SR, Taylor KD, Tremelling M, Verspaget HW, De Vos M, Wijmenga C, Wilson DC, Winkelmann J, Xavier RJ, Zeissig S, Zhang B, Zhang CK, Zhao H, Silverberg MS, Annesse V, Hakonarson H, Brant SR, Radford-Smith G, Mathew CG, Rioux JD, Schadt EE, Daly MJ, Franke A, Parkes M, Vermeire S, Barrett JC, Cho JH: Host-microbe interactions have shaped the genetic architecture of inflammatory bowel disease. *Nature* 2012, 491:119-24.
- [10] Yamazaki K, Umeno J, Takahashi A, Hirano A, Johnson TA, Kumasaka N, Morizono T, Hosono N, Kawaguchi T, Takazoe M, Yamada T, Suzuki Y, Tanaka H, Motoya S, Hosokawa M, Arimura Y, Shinomura Y, Matsui T, Matsumoto T, Iida M, Tsunoda T, Nakamura Y, Kamatani N, Kubo M: A genome-wide association study identifies 2 susceptibility Loci for Crohn's disease in a Japanese population. *Gastroenterology* 2013, 144:781-8.
- [11] Zucchelli M, Camilleri M, Andreasson AN, Bresso F, Dlugosz A, Halfvarson J, Torkvist L, Schmidt PT, Karling P, Ohlsson B, Duerr RH, Simren M, Lindberg G, Agreus L, Carlson P,

- 1 Zinsmeister AR, D'Amato M: Association of TNFSF15 polymorphism with irritable bowel
2 syndrome. *Gut* 2011, 60:1671-7.
- 3 [12] Zinovieva E, Bourgain C, Kadi A, Letourneur F, Izac B, Said-Nahal R, Lebrun N, Cagnard N,
4 Vigier A, Jacques S, Miceli-Richard C, Garchon HJ, Heath S, Charon C, Bacq D, Boland A, Zelenika
5 D, Chiocchia G, Breban M: Comprehensive linkage and association analyses identify haplotype, near
6 to the TNFSF15 gene, significantly associated with spondyloarthritis. *PLoS Genet* 2009, 5:e1000528.
- 7 [13] Nakamura M, Nishida N, Kawashima M, Aiba Y, Tanaka A, Yasunami M, Nakamura H,
8 Komori A, Nakamuta M, Zeniya M, Hashimoto E, Ohira H, Yamamoto K, Onji M, Kaneko S, Honda
9 M, Yamagiwa S, Nakao K, Ichida T, Takikawa H, Seike M, Umemura T, Ueno Y, Sakisaka S,
10 Kikuchi K, Ebinuma H, Yamashiki N, Tamura S, Sugawara Y, Mori A, Yagi S, Shirabe K, Taketomi
11 A, Arai K, Monoe K, Ichikawa T, Taniai M, Miyake Y, Kumagi T, Abe M, Yoshizawa K, Joshita S,
12 Shimoda S, Honda K, Takahashi H, Hirano K, Takeyama Y, Harada K, Migita K, Ito M, Yatsuhashi
13 H, Fukushima N, Ota H, Komatsu T, Saoshiro T, Ishida J, Kouno H, Yagura M, Kobayashi M, Muro
14 T, Masaki N, Hirata K, Watanabe Y, Nakamura Y, Shimada M, Hirashima N, Komeda T, Sugi K,
15 Koga M, Ario K, Takesaki E, Maehara Y, Uemoto S, Kokudo N, Tsubouchi H, Mizokami M,
16 Nakanuma Y, Tokunaga K, Ishibashi H: Genome-wide association study identifies TNFSF15 and
17 POU2AF1 as susceptibility loci for primary biliary cirrhosis in the Japanese population. *Am J Hum*
18 *Genet* 2012, 91:721-8.
- 19 [14] Osawa K, Takami N, Shiozawa K, Hashiramoto A, Shiozawa S: Death receptor 3 (DR3) gene
20 duplication in a chromosome region 1p36.3: gene duplication is more prevalent in rheumatoid
21 arthritis. *Genes Immun* 2004, 5:439-43.
- 22 [15] Zhang FR, Huang W, Chen SM, Sun LD, Liu H, Li Y, Cui Y, Yan XX, Yang HT, Yang RD,
23 Chu TS, Zhang C, Zhang L, Han JW, Yu GQ, Quan C, Yu YX, Zhang Z, Shi BQ, Zhang LH, Cheng
24 H, Wang CY, Lin Y, Zheng HF, Fu XA, Zuo XB, Wang Q, Long H, Sun YP, Cheng YL, Tian HQ,
25 Zhou FS, Liu HX, Lu WS, He SM, Du WL, Shen M, Jin QY, Wang Y, Low HQ, Erwin T, Yang NH,
26 Li JY, Zhao X, Jiao YL, Mao LG, Yin G, Jiang ZX, Wang XD, Yu JP, Hu ZH, Gong CH, Liu YQ,
27 Liu RY, Wang DM, Wei D, Liu JX, Cao WK, Cao HZ, Li YP, Yan WG, Wei SY, Wang KJ, Hibberd
28 ML, Yang S, Zhang XJ, Liu JJ: Genomewide association study of leprosy. *N Engl J Med* 2009,
29 361:2609-18.
- 30 [16] Bamias G, Siakavellas SI, Stamatelopoulos KS, Chryssochoou E, Papamichael C, Sfrikakis PP:
31 Circulating levels of TNF-like cytokine 1A (TL1A) and its decoy receptor 3 (DcR3) in rheumatoid
32 arthritis. *Clin Immunol* 2008, 129:249-55.
- 33 [17] Bamias G, Kaltsa G, Siakavellas SI, Papaxoinis K, Zampeli E, Michopoulos S, Zouboulis-
34 Vafiadis I, Ladas SD: High intestinal and systemic levels of decoy receptor 3 (DcR3) and its ligand
35 TL1A in active ulcerative colitis. *Clin Immunol* 2010, 137:242-9.
- 36 [18] Bamias G, Evangelou K, Vergou T, Tsimaratou K, Kaltsa G, Antoniou C, Kotsinas A, Kim S,
37 Gorgoulis V, Stratigos AJ, Sfrikakis PP: Upregulation and nuclear localization of TNF-like cytokine
38 1A (TL1A) and its receptors DR3 and DcR3 in psoriatic skin lesions. *Exp Dermatol* 2011, 20:725-31.
- 39 [19] Aiba Y, Harada K, Komori A, Ito M, Shimoda S, Nakamura H, Nagaoka S, Abiru S, Migita K,
40 Ishibashi H, Nakanuma Y, Nishida N, Kawashima M, Tokunaga K, Yatsuhashi H, Nakamura M:
41 Systemic and local expression levels of TNF-like ligand 1A and its decoy receptor 3 are increased in
42 primary biliary cirrhosis. *Liver Int* 2014, 34:679-88.
- 43 [20] Konsta M, Bamias G, Tektonidou MG, Christopoulos P, Iliopoulos A, Sfrikakis PP: Increased
44 levels of soluble TNF-like cytokine 1A in ankylosing spondylitis. *Rheumatology (Oxford)* 2013,
45 52:448-51.
- 46 [21] Meylan F, Song YJ, Fuss I, Villarreal S, Kahle E, Malm IJ, Acharya K, Ramos HL, Lo L,
47 Mentink-Kane MM, Wynn TA, Migone TS, Strober W, Siegel RM: The TNF-family cytokine TL1A
48 drives IL-13-dependent small intestinal inflammation. *Mucosal Immunol* 2011, 4:172-85.
- 49 [22] Taraban VY, Slebiada TJ, Willoughby JE, Buchan SL, James S, Sheth B, Smyth NR, Thomas
50 GJ, Wang EC, Al-Shamkhani A: Sustained TL1A expression modulates effector and regulatory T-
51 cell responses and drives intestinal goblet cell hyperplasia. *Mucosal Immunol* 2011, 4:186-96.

- 1 [23] Shih DQ, Barrett R, Zhang X, Yeager N, Koon HW, Phaosawasdi P, Song Y, Ko B, Wong MH,
2 Michelsen KS, Martins G, Pothoulakis C, Targan SR: Constitutive TL1A (TNFSF15) expression on
3 lymphoid or myeloid cells leads to mild intestinal inflammation and fibrosis. *PLoS One* 2011,
4 6:e16090.
- 5 [24] Bamias G, Mishina M, Nyce M, Ross WG, Kollias G, Rivera-Nieves J, Pizarro TT, Cominelli F:
6 Role of TL1A and its receptor DR3 in two models of chronic murine ileitis. *Proc Natl Acad Sci U S*
7 *A* 2006, 103:8441-6.
- 8 [25] Fang L, Adkins B, Deyev V, Podack ER: Essential role of TNF receptor superfamily 25
9 (TNFRSF25) in the development of allergic lung inflammation. *J Exp Med* 2008, 205:1037-48.
- 10 [26] Meylan F, Davidson TS, Kahle E, Kinder M, Acharya K, Jankovic D, Bundoc V, Hodges M,
11 Shevach EM, Keane-Myers A, Wang EC, Siegel RM: The TNF-family receptor DR3 is essential for
12 diverse T cell-mediated inflammatory diseases. *Immunity* 2008, 29:79-89.
- 13 [27] Pappu BP, Borodovsky A, Zheng TS, Yang X, Wu P, Dong X, Weng S, Browning B, Scott ML,
14 Ma L, Su L, Tian Q, Schneider P, Flavell RA, Dong C, Burkly LC: TL1A-DR3 interaction regulates
15 Th17 cell function and Th17-mediated autoimmune disease. *J Exp Med* 2008, 205:1049-62.
- 16 [28] Takedatsu H, Michelsen KS, Wei B, Landers CJ, Thomas LS, Dhall D, Braun J, Targan SR:
17 TL1A (TNFSF15) regulates the development of chronic colitis by modulating both T-helper 1 and T-
18 helper 17 activation. *Gastroenterology* 2008, 135:552-67.
- 19 [29] Schreiber TH, Wolf D, Tsai MS, Chirinos J, Deyev VV, Gonzalez L, Malek TR, Levy RB,
20 Podack ER: Therapeutic Treg expansion in mice by TNFRSF25 prevents allergic lung inflammation.
21 *J Clin Invest* 2010, 120:3629-40.
- 22 [30] Bull MJ, Williams AS, Mecklenburgh Z, Calder CJ, Twohig JP, Elford C, Evans BA, Rowley
23 TF, Slebiada TJ, Taraban VY, Al-Shamkhani A, Wang EC: The Death Receptor 3 / TNF-like protein
24 1A pathway drives adverse bone pathology in inflammatory arthritis. *J Exp Med* 2008, 205:2457-64.
- 25 [31] Calder CJ, Wang EC: An essential role for death receptor 3 in experimental autoimmune
26 uveoretinitis. *Ocul Immunol Inflamm* 2012, 20:212-4.
- 27 [32] Buchan SL, Taraban VY, Slebiada TJ, James S, Cunningham AF, Al-Shamkhani A: Death
28 receptor 3 is essential for generating optimal protective CD4(+) T-cell immunity against Salmonella.
29 *Eur J Immunol* 2012, 42:580-8.
- 30 [33] Twohig JP, Marsden M, Cuff SM, Ferdinand JR, Gallimore AM, Perks WV, Al-Shamkhani A,
31 Humphreys IR, Wang EC: The Death Receptor 3/TL1A pathway is essential for efficient
32 development of antiviral CD4+ and CD8+ T cell immunity. *FASEB J* 2012, 26:3575-86.
- 33 [34] Meylan F, Hawley ET, Barron L, Barlow JL, Penumetcha P, Pelletier M, Sciume G, Richard
34 AC, Hayes ET, Gomez-Rodriguez J, Chen X, Paul WE, Wynn TA, McKenzie AN, Siegel RM: The
35 TNF-family cytokine TL1A promotes allergic immunopathology through group 2 innate lymphoid
36 cells. *Mucosal Immunol* 2013, 7:958-68.
- 37 [35] Yu X, Pappu R, Ramirez-Carrozzi V, Ota N, Caplazi P, Zhang J, Yan D, Xu M, Lee WP,
38 Grogan JL: TNF superfamily member TL1A elicits type 2 innate lymphoid cells at mucosal barriers.
39 *Mucosal Immunol* 2013, 7:730-40.
- 40 [36] Jones GW, Stumhofer JS, Foster T, Twohig JP, Hertzog P, Topley N, Williams AS, Hunter CA,
41 Jenkins BJ, Wang EC, Jones SA: Naive and activated T cells display differential responsiveness to
42 TL1A that affects Th17 generation, maintenance, and proliferation. *FASEB J* 2011, 25:409-19.
- 43 [37] Richard AC, Tan C, Hawley ET, Gomez-Rodriguez J, Goswami R, Yang XP, Cruz AC,
44 Penumetcha P, Hayes ET, Pelletier M, Gabay O, Walsh M, Ferdinand JR, Keane-Myers A, Choi Y,
45 O'Shea JJ, Al-Shamkhani A, Kaplan MH, Gery I, Siegel RM, Meylan F: The TNF-Family Ligand
46 TL1A and Its Receptor DR3 Promote T Cell-Mediated Allergic Immunopathology by Enhancing
47 Differentiation and Pathogenicity of IL-9-Producing T Cells. *J Immunol* 2015, 194:3567-82.
- 48 [38] Cavallini C, Lovato O, Bertolaso A, Pacelli L, Zoratti E, Zanolini E, Krampera M, Zamo A,
49 Tecchio C, Cassatella MA, Pizzolo G, Scupoli MT: The TNF-family cytokine TL1A inhibits
50 proliferation of human activated B cells. *PLoS One* 2013, 8:e60136.

- [39] McLaren JE, Calder CJ, McSharry BP, Sexton K, Salter RC, Singh NN, Wilkinson GW, Wang EC, Ramji DP: The TNF-like protein 1A-death receptor 3 pathway promotes macrophage foam cell formation in vitro. *J Immunol* 2010, 184:5827-34.
- [40] Twohig JP, Roberts MI, Gavalda N, Rees-Taylor EL, Giralt A, Adams D, Brooks SP, Bull MJ, Calder CJ, Cuff S, Yong AA, Alberch J, Davies A, Dunnett SB, Tolkovsky AM, Wang EC: Age-dependent maintenance of motor control and corticostriatal innervation by death receptor 3. *J Neurosci* 2010, 30:3782-92.
- [41] Al-Lamki RS, Wang J, Tolkovsky AM, Bradley JA, Griffin JL, Thiru S, Wang EC, Bolton E, Min W, Moore P, Pober JS, Bradley JR: TL1A both promotes and protects from renal inflammation and injury. *J Am Soc Nephrol* 2008, 19:953-60.
- [42] Al-Lamki RS, Lu W, Finlay S, Twohig JP, Wang EC, Tolkovsky AM, Bradley JR: DR3 signaling protects against cisplatin nephrotoxicity mediated by TNF. *Am J Pathol* 2012, (in press).
- [43] Borysenko CW, Garcia-Palacios V, Griswold RD, Li Y, Iyer AK, Yaroslavskiy BB, Sharrow AC, Blair HC: Death receptor-3 mediates apoptosis in human osteoblasts under narrowly regulated conditions. *J Cell Physiol* 2006, 209:1021-8.
- [44] Collins FL, Williams JO, Bloom AC, Stone MD, Choy E, Wang EC, Williams AS: Death Receptor 3 (TNFRSF25) Increases Mineral Apposition by Osteoblasts and Region Specific New Bone Formation in the Axial Skeleton of Male DBA/1 Mice. *J Immunol Res* 2015, 2015:901679.
- [45] McLoughlin RM, Witowski J, Robson RL, Wilkinson TS, Hurst SM, Williams AS, Williams JD, Rose-John S, Jones SA, Topley N: Interplay between IFN-gamma and IL-6 signaling governs neutrophil trafficking and apoptosis during acute inflammation. *J Clin Invest* 2003, 112:598-607.
- [46] Hurst SM, Wilkinson TS, McLoughlin RM, Jones S, Horiuchi S, Yamamoto N, Rose-John S, Fuller GM, Topley N, Jones SA: IL-6 and its soluble receptor orchestrate a temporal switch in the pattern of leukocyte recruitment seen during acute inflammation. *Immunity* 2001, 14:705-14.
- [47] Fielding CA, Jones GW, McLoughlin RM, McLeod L, Hammond VJ, Uceda J, Williams AS, Lambie M, Foster TL, Liao CT, Rice CM, Greenhill CJ, Colmont CS, Hams E, Coles B, Kift-Morgan A, Newton Z, Craig KJ, Williams JD, Williams GT, Davies SJ, Humphreys IR, O'Donnell VB, Taylor PR, Jenkins BJ, Topley N, Jones SA: Interleukin-6 signaling drives fibrosis in unresolved inflammation. *Immunity* 2014, 40:40-50.
- [48] Wang EC, Thern A, Denzel A, Kitson J, Farrow SN, Owen MJ: DR3 regulates negative selection during thymocyte development. *Mol Cell Biol* 2001, 21:3451-61.
- [49] Stacey MA, Marsden M, Wang EC, Wilkinson GW, Humphreys IR: IL-10 restricts activation-induced death of NK cells during acute murine cytomegalovirus infection. *J Immunol* 2011, 187:2944-52.
- [50] Stacey MA, Marsden M, Pham NT, Clare S, Dolton G, Stack G, Jones E, Klenerman P, Gallimore AM, Taylor PR, Snelgrove RJ, Lawley TD, Dougan G, Benedict CA, Jones SA, Wilkinson GW, Humphreys IR: Neutrophils recruited by IL-22 in peripheral tissues function as TRAIL-dependent antiviral effectors against MCMV. *Cell Host Microbe* 2014, 15:471-83.
- [51] Wang EC, Newton Z, Hayward OA, Clark SR, Collins F, Perks WV, Singh RK, Twohig JP, Williams AS: Regulation of early cartilage destruction in inflammatory arthritis by death receptor 3. *Arthritis Rheumatol* 2014, 66:2762-72.
- [52] Strunk T, Power Coombs MR, Currie AJ, Richmond P, Golenbock DT, Stoler-Barak L, Gallington LC, Otto M, Burgner D, Levy O: TLR2 mediates recognition of live *Staphylococcus epidermidis* and clearance of bacteremia. *PLoS One* 2010, 5:e10111.
- [53] Tabeta K, Georgel P, Janssen E, Du X, Hoebe K, Crozat K, Mudd S, Shamel L, Sovath S, Goode J, Alexopoulou L, Flavell RA, Beutler B: Toll-like receptors 9 and 3 as essential components of innate immune defense against mouse cytomegalovirus infection. *Proc Natl Acad Sci U S A* 2004, 101:3516-21.
- [54] Kang YJ, Kim WJ, Bae HU, Kim DI, Park YB, Park JE, Kwon BS, Lee WH: Involvement of TL1A and DR3 in induction of pro-inflammatory cytokines and matrix metalloproteinase-9 in atherogenesis. *Cytokine* 2005, 29:229-35.

- 1 [55] Viola A, Luster AD: Chemokines and their receptors: drug targets in immunity and
2 inflammation. *Annu Rev Pharmacol Toxicol* 2008, 48:171-97.
- 3 [56] Tecchio C, Cassatella MA: Neutrophil-derived chemokines on the road to immunity. *Semin*
4 *Immunol* 2016, 28:119-28.
- 5 [57] McLoughlin RM, Jenkins BJ, Grail D, Williams AS, Fielding CA, Parker CR, Ernst M, Topley
6 N, Jones SA: IL-6 trans-signaling via STAT3 directs T cell infiltration in acute inflammation. *Proc*
7 *Natl Acad Sci U S A* 2005, 102:9589-94.
- 8 [58] Meylan F, Richard AC, Siegel RM: TL1A and DR3, a TNF family ligand-receptor pair that
9 promotes lymphocyte costimulation, mucosal hyperplasia, and autoimmune inflammation. *Immunol*
10 *Rev* 2011, 244:188-96.
- 11 [59] Slebioda TJ, Rowley TF, Ferdinand JR, Willoughby JE, Buchan SL, Taraban VY, Al-
12 Shamkhani A: Triggering of TNFRSF25 promotes CD8 T-cell responses and anti-tumor immunity.
13 *Eur J Immunol* 2011, 41:2606-11.
- 14
15





FIGURE LEGENDS










Figure 1 – DR3 expression and sources of TL1A in the peritoneal cavity. Inflammation in the cavity was induced via an i.p. injection of SES. Peritoneal membranes were harvested, fixed, sectioned and stained for DR3 expression or analyzed for the relative quantity (RQ) of TL1A mRNA by qRT-PCR. A) Representative pictures showing DR3 staining in the mesothelial layer of DR3^{+/+} mice after 0 and 12 hours but absent after 24 of inflammation (bar = 25µm.) B) Summary of DR3 expression in the mesothelial layer. Mean % positive (brown) pixels within the mesothelial layer of the membrane shown ± SEM (n=3 to 6 DR3^{+/+} mice and DR3^{-/-} mice per timepoint), ****p*<0.001 by one-way ANOVA. C) RQ of TL1A mRNA in the peritoneal membrane of DR3^{+/+} and DR3^{-/-} mice. Mean ± SEM from n=5 mice per timepoint; not significantly different by ANOVA. D) RQ of TL1A mRNA in DR3^{+/+} and DR3^{-/-} resident peritoneal macrophages 1 hour after stimulation using SES. Macrophages were sorted from other leukocytes found in the peritoneal cavity from 17 pooled DR3^{+/+} and DR3^{-/-} mice, challenged with SES and analyzed for RQ of TL1A mRNA. Mean ± SEM shown from 4 replicate experiments. **p*<0.05 by t-Test assuming unequal variance.




Figure 2 – Leukocyte subset numbers in the peritoneal cavity of DR3^{+/+} and DR3^{-/-} mice after SES and MCMV induced inflammation. Inflammation in the cavity was induced via an i.p. injection of SES or MCMV. (A) Numbers of neutrophils in the peritoneal cavity following the labeled challenges at the indicated times. (B) Peritoneal leukocytes were isolated from the cavity by lavage at the indicated times post-challenge and numbers of individual cell subsets calculated using cell counts and flow cytometry. Representative data from 1 of 2 expts. Each symbol represents mean of up to 6 mice per timepoint, error bars correspond to SEM; statistical analysis by ANOVA and Bonferroni post hoc test showed significant differences between DR3^{+/+} and DR3^{-/-} mice at ****p*<0.001, ***p*<0.01, **p*<0.05 for individual timepoints. Overall male:female ratio was 50:50 and matched for DR3^{+/+} and DR3^{-/-} groups.





Figure 3 – Chemokine production in the peritoneal cavity of DR3^{+/+} and DR3^{-/-} mice after SES-induced inflammation. Inflammation in the cavity was induced via an i.p. injection of SES. levels of the indicated chemokines were measured in: (A) cell free supernatants in peritoneal lavage by ELISA and; (B) mRNA levels in the peritoneal membrane by qRT-PCR. Representative data from 1 of 2 expts. Each symbol and error bars represent mean \pm SEM from up to 6 DR3^{+/+} and DR3^{-/-} mice per timepoint. Overall male:female ratio was 50:50 and matched for DR3^{+/+} and DR3^{-/-} groups. (C) Production of CCL3 and CCL4 by resident macrophages derived from DR3^{+/+} and DR3^{-/-} mice, stimulated with SES. Each point represents a culture; bars represent mean \pm SEM. Values below the limit of the assay were assigned the lowest limit for generation of means and statistical testing. Dotted line shows lowest limit of assay. Statistical analysis by ANOVA and Bonferroni post hoc test showed significant differences between DR3^{+/+} and DR3^{-/-} mice at *** p <0.001, ** p <0.01, * p <0.05.

Figure 4 – Thickness of the mesothelial layer and collagen deposition in DR3^{+/+} and DR3^{-/-} mice after inflammation induced by multiple applications of SES. Fibrosis of the mesothelial layer was induced by 4 i.p. injections of SES. Representative micrographs of the peritoneal membranes visualized with (A) hematoxylin and eosin, and (B) van Gieson stain, from mice treated as indicated. M = mesothelial layer; Bar = 50 μ m. Summary of (C) mesothelial thickness and (D) collagen deposition. Bar and error bar represents mean \pm SEM from up to 6 mice. Statistical analysis by ANOVA showed significant differences between DR3^{+/+} and DR3^{-/-} mice *** p <0.001 by Bonferroni post hoc test. Overall male:female ratio was 50:50 and matched for DR3^{+/+} and DR3^{-/-} groups.









Supplemental Table 1 – Counts for leukocyte subsets are not different in the unchallenged peritoneal cavity or peripheral blood of DR3^{-/-} and DR3^{+/+} mice

Cell subset	Peritoneal cavity *			Peripheral blood No/ml †		
	DR3 ^{+/+}	DR3 ^{-/-}	Significance ‡	DR3 ^{+/+}	DR3 ^{-/-}	Significance ‡
Macrophages/Monocytes	7.2 ± 1.8 x10 ⁵	6.4 ± 1.6 x10 ⁵	NS	4.2 ± 1.0 x10 ⁵	4.7 ± 1.6 x10 ⁵	NS
Dendritic cells	5.1 ± 1.6 x10 ³	3.3 ± 0.3 x10 ³	NS	N/A [§]	N/A [§]	N/A [§]
Neutrophils	N/A [§]	N/A [§]	N/A [§]	2.3 ± 0.6 x10 ⁵	4.1 ± 2.2 x10 ⁵	NS
Eosinophils	1.9 ± 0.7 x10 ⁵	7.6 ± 0.6 x10 ⁴	NS	5.2 ± 1.3 x10 ⁴	3.5 ± 0.8 x10 ⁴	NS
T Cells	4.5 ± 0.5 x10 ⁵	4.6 ± 1.1 x10 ⁵	NS	4.0 ± 0.5 x10 ⁵	4.5 ± 0.6 x10 ⁵	NS
B Cells	8.6 ± 1.0 x10 ⁵	9.3 ± 2.9 x10 ⁵	NS	9.6 ± 1.8 x10 ⁵	1.0 ± 0.2 x10 ⁶	NS
NK Cells	6.8 ± 1.7 x10 ⁴	4.7 ± 0.8 x10 ⁴	NS	8.3 ± 1.4 x10 ⁴	7.2 ± 1.0 x10 ⁴	NS
NKT Cells	7.2 ± 1.8 x10 ⁴	6.9 ± 0.6 x10 ⁴	NS	1.3 ± 0.2 x10 ⁴	1.3 ± 0.2 x10 ⁴	NS

* Values correspond to number of peritoneal leukocytes, mean ± SEM (n=5)

† Values correspond to number of peripheral blood leukocytes/ml, mean ± SEM (n= 4 DR3^{+/+} or 5 DR3^{-/-})

‡ Significance between the two genotypes using t-Test, NS = no significant difference

§ N/A corresponds to subsets either not present or not tested




Supplemental Table 2 – Leukocyte proliferation in the peritoneal cavity following SES challenge

Cell Subset *	Proportion of proliferating cells [†]		
	DR3 ^{+/+}	DR3 ^{-/-}	Significance [‡]
Resident macrophages	42.4 ± 13.2	44.5 ± 8.2	N.S.D
Inflammatory macrophages	43.7 ± 11.5	47.9 ± 7.7	N.S.D
T Cells	49.3 ± 2.3	50.5 ± 7.5	N.S.D
B Cells	36.9 ± 2.8	28.8 ± 8.2	N.S.D
NK Cells	35.2 ± 10.4	38.2 ± 3.8	N.S.D
NKT Cells	47.7 ± 3.8	51.8 ± 10.1	N.S.D

* Ki67⁺ cells were assessed in the indicated cell populations using the following phenotypic criteria: resident macrophages (F4/80⁺CD11b⁺); inflammatory macrophages (F4/80^{int}CD11b^{int}); NK cells (NK1.1⁺αβTCR⁻); NKT cells (NK1.1⁺αβTCR⁺); T cells (CD3⁺αβTCR⁺NK1.1⁻); B cells (CD19⁺B220⁺)

[†] Values = %Ki67⁺ of each subset 48 hours after SES challenge (mean ± SEM; n= 5 DR3^{+/+} or 6 DR3^{-/-}). Gating as indicated in Supplemental Figure 1.

[‡] Mann Whitney test showed N.S.D = no significant difference between DR3^{+/+} and DR3^{-/-}.






Figure Legends

Figure 1 – Flow cytometric gating for determining Ki67⁺ proliferating cells. Mice were challenged with SES and peritoneal leukocytes isolated from the cavity by lavage, stained for cell subsets, permeabilised and proliferation measured using the intracellular marker Ki67 and DAPI to differentiate binucleate non-proliferating cells. Representative plots of total peritoneal leukocytes showing isotype control and KI67 staining from DR3^{+/+} and DR3^{-/-} mice. Early proliferative G1 cells (Ki67⁺ DAPI^{lo}), S and G₂M phase (Ki67⁺ DAPI^{int/hi}) and non-proliferating binucleate cells (Ki67⁻ DAPI^{hi}) shown.

Figure 2 - Chemokine levels in the naïve peritoneal cavity are not different between DR3^{+/+} and DR3^{-/-} mice. The peritoneal cavity was lavaged, cells spun down and chemokine concentrations in the supernatant measured by ELISA. Data from n=3 DR3^{+/+} and DR3^{-/-} mice where available, but some values were below the manufacturer's detection threshold of the ELISA and therefore excluded from analysis. No significant differences were detected.